



ALLEN SWCD 2025 FINGERLING FISH SALES




Bluegills are popular food for largemouth bass and humans. Most bluegills can spawn by age two when they are about 3 inches long. Young bluegills feed on tiny, microscopic plants and animals. Adults feed on insects, snails, small crayfish, and fish eggs. Bluegills can grow up to six inches in two to four years if provided enough food and space, and some have even grown over 10".




Hybrid Bluegill are crossed between a male bluegill and female green sunfish. This cross produces a fish with the large mouth of the sunfish and the aggressiveness of the bluegill, which enables the fish to grow faster and larger than the common breed. It will feed on a commercial feed with annual growth rates of up to ½ - ¾ pounds per season. These fish can reach weights of 2-2 ½ pounds.



Largemouth Bass are the number one predator in ponds. They begin spawning about the age of three when they are 10–12" long. Adults' diets consist of insects, crawfish, and small fish (bluegill and small largemouth bass). Largemouth bass grow to an average size of 1–2 pounds and live for 4-6 years.



Red Ear Shell Cracker thrives in clear, warm impoundments with rooted vegetation. They are attracted to submerged vegetation and trees. In well-managed waters with good habitats, red-ear sunfish will reach 9-11" in length



Channel Catfish are mostly bottom feeders, preferring live or dead insects, crayfish, and occasionally aquatic plants. Catfish tend to stir up the bottoms of shallow ponds and cause the water to become muddy. Their average adult size is 14-16".



White Amur are members of the minnow family. They usually live 10 to 15 years and can grow to weigh 100 pounds. White Amur feeds on leafy, rooted aquatic plants, such as pondweed and coontail. Lacking these, they will consume floating duckweed, green algae, and even cattails.



Perch prey upon aquatic insects, invertebrates, a wide variety of fishes, and the eggs of other fish species. Adults average between 7-12" and weigh between 4 ounces and 1 pound.

Black Crappie are part of the sunfish family. They spawn in late spring, the same time as bluegill, and primarily feed on insects and smaller fish. They are best caught in the colder months and are a favorite of ice fishermen. In the right conditions, they can grow to eleven or twelve inches.

Japanese Koi is a domesticated version of the common carp. These Fish come in many colors and varieties and are a great addition to your backyard pond.

Minnows are small fish that swim in schools and are used as feeder fish. We recommend 1000 per surface acre to allow for breeding.

ALL FISH ARE LIVE AND MUST BE PICKED UP IN PERSON